

TR-01-342

2427

ARL-TR-65-15

STUDY OF MONKEY, APE, AND HUMAN MORPHOLOGY AND PHYSIOLOGY
RELATING TO STRENGTH AND ENDURANCE

PHASE IX

THE STRENGTH TESTING OF FIVE CHIMPANZEE AND SEVEN HUMAN SUBJECTS

William E. Edwards
Thomas Erskine Clarke

August 1965

SEP 15 1965

DDG-IRA E

6571st Aeromedical Research Laboratory
Aerospace Medical Division
Air Force Systems Command
Holloman Air Force Base, New Mexico

2427
20090505 277

2427

The animals used in this study were handled in accordance with the "Principles of Laboratory Animal Care" established by the National Society for Medical Research.

This report may be reproduced to satisfy needs of U.S. Government agencies. No other reproduction is authorized except with permission of the 6571st Aeromedical Research Laboratory, Holloman AFB, NMex.

This report is made available for study with the understanding that proprietary interests in and relating thereto will not be impaired. In case of apparent conflict or any other questions between the Government's rights and those of others, notify the Judge Advocate, Air Force Systems Command, Andrews Air Force Base, Washington, D.C. 20331.

Certain of the animals used for this research were provided by the National Aeronautics and Space Administration under Order R-25.

Do not return this copy. Retain or destroy.

AD-469585

STUDY OF MONKEY, APE, AND HUMAN MORPHOLOGY AND PHYSIOLOGY
RELATING TO STRENGTH AND ENDURANCE

PHASE IX

THE STRENGTH TESTING OF FIVE CHIMPANZEE AND SEVEN HUMAN SUBJECTS

William E. Edwards
Thomas Erskine Clarke

FOREWORD

This is the last in a series of nine papers concerned with the Study of Monkey, Ape, and Human Morphology and Physiology Relating to Strength and Endurance. This study was conducted in part under Contract AF 29(600)-3466, Project 6892, Task 689201, monitored by Major James E. Cook, Veterinary Services Division, ARV.

In 1958 the writer initiated a study on the strength of humans and other primates. This study was conducted from 1959 to 1961 at the University of Chicago as a National Science Foundation Science Faculty fellow. In 1961 the author was employed as a consultant by the 6571st Aeromedical Research Laboratory and the co-author of this paper was employed as his assistant to conduct a program of training and testing chimpanzees for flexion strength of the arms. The data acquired was analyzed in 1963 and compiled for this report.

The author wishes to acknowledge the cooperation of Lt Col Frederick H. Rohles, Lt Col Hamilton H. Blackshear, Major Clyde H. Kratochvil, Major James E. Cook, Major Marvin E. Grunzke and Major Robert H. Edwards, all of the 6571st Aeromedical Research Laboratory for their help in the preparation of this paper.

Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

C. H. Kratochvil

C.H. KRATOCHVIL

Major, USAF, MC

Commander

ABSTRACT

Five chimpanzees (two immature of each sex and one adult male) were trained for testing of elbow flexion strength. Resulting scores were compared with those of seven young (20 to 37 years) adult human males. The apes manifested a 2-1/2 fold superiority by body-weight and, sex and age equivalent, an appreciable superiority by brachial cross-sectional area.

TABLE OF CONTENTS

	Page
1. INTRODUCTION	1
2. RESEARCH HISTORY	1
3. RESEARCH METHODOLOGY	2
4. DESCRIPTION OF HUMAN SUBJECTS	2
5. EXPERIMENTAL RESULTS OF HUMAN STRENGTH TESTING	6
6. METHODOLOGY OF TRAINING AND TESTING CHIMPANZEES	8
7. DESCRIPTION OF CHIMPANZEE SUBJECTS	8
8. EXPERIMENTAL RESULTS OF CHIMPANZEE STRENGTH TESTING	10
9. CONCLUSIONS	15
REFERENCES	21

LIST OF ILLUSTRATIONS

Figure

1. Assistant Author, Fourth Chimpanzee, and Testing Apparatus	3
2. Fourth Chimpanzee at Start of Pull	4
3. Fourth Chimpanzee in Early Stage of Successful Lift	5

1. INTRODUCTION

Various types of anecdotal and other general observational evidence have indicated that chimpanzees are very much stronger than humans. Some tests (Bauman, 1923 and 1926) have corroborated such indications of strength superiority, while others (Finch, 1943) have indicated no appreciable difference.

Knowledge of whether marked differences in strength exist in chimpanzees and humans is highly significant to relevant aspects of pure science and of applied science, such as to the development of the man-in-space program (Edwards, 1965b, pp. 1-2) -- for example, marked differences in muscular strength are associated with appreciable differences in tissue damage under high acceleration forces.

The training and testing program herein reported was effected to determine valid strength comparisons between the two primates.

2. RESEARCH HISTORY

In 1959, the writer conducted fairly extensive library research on the problem of relating the strength of man and other primates to size, at which time it was recognized that the most ideal non-human subjects were chimpanzees -- because the only significant prior studies of non-human strength had utilized the chimpanzee, because of their better responsiveness to training in some respects, and especially because their comparisons with humans would be facilitated by their approximately equivalent body-size. Derived data on the chimpanzee would also be more directly applicable to certain practical problems -- including some of those involved in space flight, as already noted.

The senior author was employed by the 6571st Aeromedical Research Laboratory as a consultant in 1961 and the results of this investigation were subsequently prepared for this report under Air Force Contract 29(600)-3466.

The author and the co-author collaborated in the training and testing program for this study.

3. RESEARCH METHODOLOGY

On a steel platform 128.6 by 71 cm. was mounted a steel chair with adjustable seat and back as well as adjustable neck, chest, thigh, and ankle braces to restrain the simian subjects of the experimentation (Figs. 1-3). An adjustable 53.6 by 45.6 cm. steel table was affixed immediately in front of the subject. Upon this table was mounted a handle of 3.7 cm. (almost 1.5 inches) diameter, readily shifted to either right or left side, attached perpendicularly to a rod which slides within a cylinder, the angle of which to the table was adjustable. Attached in turn to the handle-rod was a 0.39-cm.-diameter, seven-strand twisted steel cable, which connected via a system of pulleys to a rectanguloid steel bucket with a capacity of more than 400 pounds of measured weights. Alternatively, the cable was attached to a dynamometer, but testing revealed proportionately large inaccuracies and caused it to be abandoned; all of the scores here reported are the actual weights of bucket plus weights. The handle-rod in its cylinder, the cable, and the pulleys were all well-fitted and well-lubricated, so the Holloman Air Force Base machinists who meticulously fashioned the apparatus were confident that friction was reduced to less than 1 per cent, in contrast to the enormous friction of the apparatus of others, such as that of Finch (1943).

The bucket was kept either suspended entirely so that the slightest lift would be evident or was in contact with the floor only along one edge. The latter arrangement had the psychological advantage of encouraging the subject by permitting slight movement of the handle before the maximum force required was applied, while simultaneously insuring that no lift would be incorrectly recorded as successful since complete lifting of the bucket along the contacting edge was quite evident when it occurred.

It was possible to satisfy virtually all of the thirty-seven criteria previously formulated by the writer as essential to proper comparisons of ape and human strength (Edwards, 1963a). Among these criteria, the number of muscles involved in the given performance is especially crucial; the relatively large forces which can be developed but minimal number of muscles which contribute to the force make elbow flexion perhaps as near-ideal for this purpose as is possible (Wilkie, 1950, pp. 250-251 and 266-269).

4. DESCRIPTION OF HUMAN SUBJECTS

The seven human subjects were all adult males in good health and with perhaps one exception at least somewhat active in physical sports. The three heaviest subjects had the highest proportion of adipose tissue, with the third lightest intermediate in this respect between the three heaviest and the remainder. Although he manifested some excess adipose tissue, the heaviest subject was quite muscular despite his weight, largely through having just participated for one year (June, 1960, to



Figure 1
Assistant Author, Fourth Chimpanzee, and Testing Apparatus




Figure 2

Fourth Chimpanzee at Start of Pull, Left Hand and Overhanded
(When turned on, the light indicates a lift should
be attempted for positive and negative incentives.)




Figure 3

Fourth Chimpanzee Momentarily after Near-Maximal Weight
Has Cleared Floor in Early Stage of Successful Lift

May, 1961) in a standard weight-lifting program to which he devoted one hour per day. The third and sixth subjects had years earlier scored in approximately the top two percentiles in "physical fitness index" tests -- primarily of strength and secondarily of endurance (Cureton, 1947). In summary, however weaker than their physically much more active distant and even very near ancestors, the majority of the human subjects employed in this study were quite certainly at least moderately stronger than average Americans, sex, size, and age considered.

All subjects but the lightest -- who was tested a day later -- were tested 4 September, 1961.

Data on the subjects, all of whom except the fifth are right-handed, are provided in Table 1.

5. EXPERIMENTAL RESULTS OF HUMAN STRENGTH TESTING

The training of human subjects consisted of verbal instruction and of brief trials to provide sufficient familiarity with the apparatus for near-optimum functioning during the tests; since the performance required only very simple patterns of muscular contraction (a major consideration in the design of the apparatus), procedural familiarity quite surely did not decrease maximal scores at all appreciably (Cureton, 1947), and any slight decrease due to this factor was at least fairly equivalent for all the human subjects.

Motivation consisted of a combination of interest in and desire to aid the research project, curiosity regarding the subject's own strength, and the competitive desire to score favorably in comparison with others. A given performance was generally observed by several of those selected for testing, and all to some degree sought and, in an attempt by the writer to maximize all of the human scores, were informed of the scores of the others.

To minimize the influence of the fatigue factor, the subjects were given rest-intervals between individual pulls until they reported feeling rested. At near-maximal pulls, these rest-intervals averaged slightly more than two minutes. Between series (A, B, C, and D), the rest-intervals approximated five minutes while adjustments were made. To provide indications of the fatigue factor, the sequences of right versus left and underhanded versus overhanded series were varied and recorded; comparisons reveal clearly that the subjective appraisals of essentially complete recovery from fatigue were probably erroneous in every case.

Rough approximations of the optimum angles were determined by brief experimentation prior to the series here recorded. For the testing of all overhanded pulls, the handle-rod was set at horizontality. In preparation for overhanded pulls, the table was adjusted to the height and distance from the subject which, at the moment the raising of the

Table 1

Anthropometric Data on the Human Subjects
(in centimeters or square centimeters where not otherwise indicated)

Characteristic	Subject						
	1	2	3	4	5	6	7
Age (in years)	30	20	33	36	37	35	28
Weight (in pounds)	125.5	144.5	167.0	169.0	186.5	221.0	248.5
Stature (in inches)	68.0	71.5	69.0	68.7	71.4	74.7	73.0
Mid-brachial girth*							
Right	24.0	27.6	30.0	28.4	30.8	34.6	35.8
Left	23.2	26.5	29.4	27.2	32.6	32.2	35.3
Distance of maximum forearm girth from proximal tip of ulna (olecranon)							
Right	4	6	5	6	8	4	8
Left	4	6	5	6	8	4	8
Maximum forearm girth*							
Right	25.4	26.2	27.5	27.3	27.3	30.1	35.1
Left	25.0	26.1	27.2	26.5	27.8	29.3	34.9
Distance of minimum forearm girth above distal tip of ulna (styloid process)							
Right	2.5	3	2.5	3	3	3	1.5
Left	2.5	3	2.5	3	3	3	1.5
Minimum forearm girth*							
Right	15.9	16.2	17.2	15.9	17.2	19.1	28.8
Left	15.6	15.7	16.4	15.7	17.7	18.6	28.8
Distance from acromion to tip of ulna (styloid process), measured palm down							
Right	56.6	57.1	53.6	55.4	57.0	60.4	58.2
Left	56.4	57.5	52.6	54.9	56.8	60.0	59.0
Ulna length (olecranon to styloid process)							
Right	27.8	29.1	26.4	27.4	30.3	29.2	29.5
Left	27.4	28.5	26.0	26.6	30.0	28.4	29.7
Brachial cross-sectional area**	44.39	58.21	70.17	61.48	79.94	88.75	100.54

*Girths were measured with the limb extended and relaxed

**Calculated from average mid-brachial girth on assumption of round section, which slightly overstates actual cross-sectional area

weight commenced, provided an angle of the forearm's main axis sloping down toward the subject of 5 degrees from the horizontal and of the upper-arm's main axis sloping up toward the subject at 6 degrees. For under-handed pulls, the handle-rod sloped up toward the subject at 10.5 degrees, while the forearm and upper arm angles were increased to 18 and 23 degrees respectively. These angles were approximated as closely as was feasible for all subjects, except for underhanded series D2 (immediately following D1) for the second subject, for which the slightly lower maximum score was obtained with the handle-rod horizontal.

To insure insofar as was feasible that the muscles tested were those providing flexion of the forearm at the elbow-joint, in all but one brief series the shoulder of the subject was braced by the arms of assistants and by boards held by them to minimize shoulder movement, and the changing angle of flexion at the elbow was observed while the handle was drawn back (on all but a few of the successful lifts to its maximum range of motion of approximately 3.5 cm.). It should be emphasized that in the one series designed to indicate the approximate degree of superiority of scapular muscles -- D2 for the third subject, immediately following D1 -- the upper extremity was held rigidly straight at the elbow, so no effort was required of the flexors at the elbow to maintain flexion at that joint.

The results of the tests are recorded in Table 2.

6. METHODOLOGY OF TRAINING AND TESTING CHIMPANZEES

The high interindividual and even intraindividual variability in chimpanzee characteristics such as temperament, cooperativeness, and response to electric shock can hardly be overemphasized. For this reason, a wide variety of both positive and negative motivations, which need not be discussed here since they have already been considered in another recent paper (Edwards, 1963a), were employed. By the somewhat unorthodox adjusting of motivation to fit the individual subject, it was possible thoroughly to train the chimpanzees -- none of which had experience with trained performance requiring maximum muscular effort -- with great rapidity.

The primary test of adequate training for strength testing is whether maximum scores in different series closely approximate one another, for such correlations should occur if consistent and adequate motivations are operative and if the limiting asymptote of muscular strength is fairly closely approached (Edwards, 1963a). The data presented subsequently in this paper demonstrate the adequacy of this training.

7. DESCRIPTION OF CHIMPANZEE SUBJECTS

Large quantities of panometric and other types of data on the chimpanzee subjects of the experimental program were recorded. Only the data

Table 2
Weightlifting Performances of Human Subjects

Subject and Series Sequence	Lifts Attempted	Maximum Lift	Average Maximum Lift	Body-Weight (in pounds)	Average Maximum Lift in Pounds per Pound of Body-Weight	Average Brachial Cross-Sectional Area	Average Maximum Lift in Pounds per Square Centimeter
1(C,D,B,A)*	A 80,100,120,140,(160)**(150)	140	122.5	125.5	.976	44.39	2.76
	B 80,100,120,(140),(130),(130)	120					
	C 80,100,120,140,(145),(145)	140					
	D 80,(100),(90),90,(95)	90					
2(C,D,A,B)	A 100,120,140,150,160,180,200,190,200,(205)	200	161.3	134.4	1.108	58.21	2.77
	B 80,100,120,(130),125,(130)	125					
	C 80,100,100,120,140,160,180,200,(210)	200					
	D1 100,120,(140),(130),(120),(120)	120					
	D2 (120),100,(100),110						
3(A,B,C,D)	A 80,100,120,140,160,180,(185),185,(190),(190)	185	156.3	167.0	.936	70.17	2.23
	B 80,100,120,(140),135	135					
	C 80,100,120,140,(160),(160),150,160,(170),165,(170)	165					
	D1 80,100,120,130,140,(150),(145)	140					
	D2 155,165,(170)						
4(B,A,C,D)	A 130,150,(180),(160),150,155,(160)	155	150.0	169.0	.888	61.48	2.44
	B 100,120,130,140,(150),(140),130,135,140,145,(150),(150)	145					
	C 100,150,120,170,(180),(175),175,180,(190),(190),(185)	180					
	D 80,100,(120),120,(125),(125)	120					
5(D,C,A,B)	A 80,100,120,140,150,160,(170),(170),(165)	160	136.3	186.5	.731	79.94	1.70
	B 80,100,(120),(110)	100					
	C 80,100,120,140,(150),150,155,(160)	155					
	D 80,100,120,130,(140)	130					
6(A,B,D,C)	A 80,100,120,140,160,180,200,220,225,(230),230,(235)	230	187.5	221.0	.848	88.75	2.11
	B 80,100,120,140,150,155,160,(165)	160					
	C 80,100,120,140,160,180,200,210,220,(230),(225)	220					
	D 80,100,120,140,(150),(145)	140					
7(A,B,D,C)	A 110,130,150,170,190,190,210,230,250,(260),(255)	250	200.0	248.5	.805	100.54	1.99
	B 80,100,120,140,150,160,(180),170,(175),(175)	170					
	C 110,130,150,170,190,210,230,240,(250),(250)	240					
	D 80,100,120,140,(160),(150),140,(145)	140					

A = Right, overhanded
B = Right, underhanded
C = Left, overhanded
D = Left, underhanded

*Sequence of the four tested

**parentheses enclosing a score denote strenuous attempt but failure to lift that weight

most relevant to comparisons of chimpanzee and human size and strength are given in Table 3.

8. EXPERIMENTAL RESULTS OF CHIMPANZEE STRENGTH TESTING

The authors wrote as many as 34 pages of condensed notes on cards per day of the testing program; at night such data were even more extensively recorded on "dictaphone" belts, which in toto constitute the equivalent of several hundred typed pages. It is therefore evident that high selectivity must be utilized in the succinct presentation of experimental results.

Such data are presented in highly condensed form in the following pages. All successful scores and all unsuccessful attempts (in parentheses) involving near-maximal effort are listed in sequence for every day each of the five subjects was tested; attempts for which a marked effort was not required or at least was not given are not here recorded. A semicolon represents a major temporal hiatus in the testing, generally with appreciable resting by the subject. In the data listed below for each of the five subjects successfully trained, each lift recorded in a series was attempted with the arm, right or left, indicated at the start of the series. Occasionally the subject changed from an overhanded to an underhanded grip or vice versa during a series; the grip for each attempt was that indicated at the start of the series except where individually indicated to the contrary in the list.

Prior to 25 August, 1961, tests were conducted in a slightly simpler steel restraining chair than that subsequently employed; the weight was attached to a rope which, after passing around a pulley, was in turn attached to a simple rod handle of wood. All tests on and after 25 August were conducted with the more elaborate steel apparatus previously described.

To the data which follow may be added the fact that only the largest chimpanzee (which, despite his apparent viciousness, proved to be the most cooperative of all) was anaesthetized (by sodium pentathol, followed by the counteracting drug mikedimide) before being moved from his cage to the restraining chair -- and even he returned from the testing building without restraint. But the smaller subjects were unwilling participants in the program, and as many as six skilled handlers required as much as 45 minutes to subdue and force one of them into the restraining chair.

It should also be observed that fairly extensive attempts were made on the two most tractable subjects, the third and the fifth, to test differences in muscular force at varying angles of elbow flexion. But in the time available, the two subjects could not learn to pull at markedly different (and apparently somewhat less "natural") angles; the fundamental change required was apparently never comprehended by the apes.

Table 3

Panometric Data on the Chimpanzee Subjects
(in centimeters or square centimeters where not otherwise indicated)

<u>Characteristic</u>	<u>Subject</u>				
	1	2	3	4	5
Sex	Male	Female	Female	Male	Male
Age (in months)	65	56(est.)	54(est.)	68(est.)	144(est.)
Weight (in pounds)	45.5	48.0	56.0	60.0	98.0
Maximum forearm girth*					
Right	22.5	23.0	22.6	23.2	30.5
Left	22.7	22.0	22.9	24.3	29.8
Mid-brachial girth*					
Right	21.8	22.0	20.6	21.5	30.5
Left	21.9	22.0	21.5	22.5	30.1
Brachial cross-sectional area**	37.99	38.51	35.27	38.51	73.06

*Girths were measured with the limb extended and relaxed

**Calculated from average mid-brachial girth on assumption of round section, which slightly overstates actual cross-sectional area

Subject 1 (ARL #80)

8/25 (right, overhanded): 80, 90, (100), (100), 80, 90, 100, 105; (105); (105), 80, 90, 100, (103), 110; (112), (112), (112), 112, (112), (112), (100), 80, 90, 100, 110, 112, 112, 116, (118), (118), (118); (118), (117), 80, 100, 110, (115), 115, (116), (116), 116, 117, (120), (120), (120), (120), (120), (120).

8/29 (left, overhanded; while recuperating from illness): (70), (70), 70, 70, (70), (70), 70, 75, 60, 70.

9/6 (left, overhanded): 80, 100, 110, 120, 130, 140, (145), (145), (145).

Subject 2 (ARL #46)

8/7 (right, overhanded): 74, 79, 84, 89, 94, 100, (105).

8/8 (right, overhanded): 74, 101, (109), (109), (109), (109), (109; weight almost completely lifted for total of 5 seconds of 7-second interval encompassing last two attempts).

8/21 (right, overhanded): 90, 90, 100, 105, 110, (115), (115).

8/22 (right, overhanded): 70, 80.

8/24 (right, overhanded): 80, 80, 90, 100, (105), (105); 80, 90; 80; 80, (90), (90); 80, 90, 100, 103, 110 (both hands), (110), (110), 110, (112), (112; under), 100, (110); 80, 80, 80, 90, (100), 100, 103, 110, (112), (112), (112); 100, 100.

8/24 (right, underhanded): 80, 80, 90, 90, (100), (100), 100, 100, 100.

Subject 3 (ARL #35)

8/19 (left, overhanded): 90, 100, 110, (110), 110, (120); 90, 110.

8/29 (left, overhanded): 80, 80, 90; 80, 90, 100; 80, 80, 90, 100, 105; 90, 100, 105, 108, 110, 112, 100, 110, 115.

8/30 (left, overhanded): 80, 90, 100, 105, 110, 113, 115, 118, (120), (120), (120), (120), 120, (122), (122); 100, 110, 115, 120, 122, 125, 127, (127), (127), (127); 80, 80, 90, 100, 110, 115, (120), 120, 125, 127, 130, (132), (132), (132), (132), (132).

8/30 (right, overhanded): 80, (90), (90), 80, 90; 80, (90), 80, (90), 80, 80, 80, 80, 80, 90, (100).

8/31 (left, overhanded): 80, 90, 100, 110, 120, 125, 130, (132), (132), (130).

8/31 (right, overhanded): 80, 100, 105; 80, 90; 80, 90; 80, 80.

9/2 (right, overhanded): 80, 90, 100, 110; 80; 80, 85, 90, 100; 80, 90, (110), (110), 100; 80, 90; 80, 90, (95), (95), (95).

9/6 (right, overhanded): 2 hours of attempted training and testing at several handle-shaft, brachial, and antebrachial angles; maximum lift only 60.

Subject 4 (ARL #32)

8/21 (right, overhanded): 80; 50.

8/22 (right, overhanded): 80, 90, (100), 90, 100, 105, (110); 80, 90, 100, 105, 110, 110, (115), (115); 80, 80, 90, 90, 100, 100, 110, (115), (115), (115).

8/23 (right, overhanded): 80, 100, 100, (115), (115), (115), (113), 110, (111); 80, 90, 100, (110); 80, 90; 80, 80, 100, 110, (115), (115), (115), (115).

8/23 (left, overhanded): 70, 70, 90, 90, 100, 105, 110, (115), 115, (120), (120), (117), (117), (117), (117), 117, (120).

8/26 (right, overhanded): 80, 90, 100, 105, 110, 112, 115, 118, (120), (120), (120), (120), (120), (120), (120); 80, 100, 110, (115), (115), (115), (115), 110, 113, 115, 118, (120), (120), (120), (120).

8/26 (right, underhanded): 70, (90), 80, (90), (90), (90), (80).

8/26 (left, overhanded): 80, 90, 105, 115, 120, 125, 127, 125, 127, 130, 135, (140); 130, (140), (140), (140; under); 100, 110; 100, 100, 110, 120, (130), 120, 130, (135), (130), (130); 80, 80, 80, 100, 115, 125, (130), (130).

8/26 (left, underhanded): 30.

8/30 (right, overhanded): 80, 100; 80.

8/31 (right, overhanded): 80, 80, 90, 90; 80, 80, 90; 80, 90, 90, 95, 100, 105; 90; 80, 90, 100; 80, 80, 80, 90, 100, 105, 105, 110; 90; 80, 80, 90, 100, 105 (under), 105; 80, 80, 90, 100, 105, 107, 110, 112, (115); 80, 80, 90, 100, 110, 115, 120, 122, 127, 130, 135 (barely lifted).

9/2 (right, overhanded): 80, 90, 100, 110, 115, 117, 120, 125, (127), 127, 130, (132), (132), (132); 80, 80, 80, 80, 100, 110; 80.

9/2 (right, underhanded): 80, 90, (100), 95, 100, (105), (105), 90 (over), 90, (100), 100, (105), 105, 110, (115), 115, (120), (120), (120), (120), (120).

9/2 (left, overhanded): 80, (90; under), 90, 100; 90, 100, 110, (120), 120, 125, (130), 130, 135, (138), (138).

9/2 (left, underhanded): 80, 100, 110 (over).

Subject 5 (ARL #134)

8/28 (right, overhanded): 180, 230, 200, 200, (250), 250, (280), (280), 180, 230, (250), (250), 250, (260), (260), (260), (260), (260), (260), (260), 260; 230, 250, (260), 260 (under), (270; under), 270, (280), (280; under), (280), (280), (280; under), (280; under), (280), (280), (280); 230, 230, 230, 230; 180, 230, 250 (under), (250), (250; under), (250), (250; under), (250), 250 (under), (230), 230 (under).

8/28 (left, overhanded): 180, (210), (210), 210, 210, 230, 240, 240, 250, 260.

8/28 (both, overhanded): 270, (270; left under), 270, 270 (writer slapped subject's hand and verbally indicated disapproval), 270 (subject raised digit III of right hand and showed to writer, apparently for approval of alternative plan to lift weight, unsuccessfully attempted by left arm only and underhanded 4 minutes earlier; not understanding, writer replied "fine"; subject immediately placed this digit overhanded on handle beside overhanded left hand and readily lifted weight), 270, 270, 270, 250, 250, 250, 250, 250, 250, 250, 250, 230 (left, over), 250 (left, under), 260 (left, over, plus digit III of right hand), 300 (left, over, plus digits II and III of right hand), 330 (left, over, plus digits II-IV of right hand), (380; steel platform of apparatus bent, so unable to raise weight), 380 (left, over, plus digits II and III of right, under; back of platform weighted, so bent less), (410; platform bent barely too much for weight to rise).

9/1 (right, overhanded): 180, 230 (under), 230; 230; 130, 130, 180, 230 (under), 250 (under), (260; under), (260), 250 (under), (260; under), (260), (260; under), (260), (255), (255; under), (255; under), (230; under), 230, 230 (under), 230, (250), (255), (255), (255), (255), (255; under), (255), (255), (255; under), (255), (250), (250), (250; under), (250; under), (250); 180, 230, 250, (260), (260), (260), 260, (260), 265.

9/5 (right, overhanded and underhanded): 7 hours of attempted training and testing at varying handle-shaft heights and angles and at varying brachial and antebrachial angles; apparently no pulls in alignment with the handle-shaft; despite repeated effort by subject, maximum lift only 160.

The results of all of the strength testing of chimpanzees are summarized in Table 4.

Scores lacking clear indication of being normal maximal scores were omitted or set in parentheses, and were in either case excluded from subsequent calculations. The final scores of the first subject were the only scores excluded as excessive. This subject had on 6 September almost escaped after being placed in the restraining chair, and resubduing required half an hour of great effort, so it is felt that these scores were probably raised "abnormally" by epinephrine release and associated phenomena. Similar phenomena -- in addition to a moderate degree of muscular hypertrophy resulting from the training and testing program -- likely affected the scores of the fourth subject on 2 September, for the tests on that date were associated with frequent screaming and struggling, but the presumed increase by epinephrine release was apparently counter-balanced by fatigue. For reasons obvious upon inspection of the data previously presented, other scores were not included in calculations because they were quite surely markedly less than potential maxima. The percentages by which the underhanded lifts were less than the over-handed lifts for the three subjects for which such scores were available were calculated -- second, 9.1; fourth, 14.8; and fifth, 3.7 and 3.8. With scores for both arms, the fifth subject determined half the average difference of 7.85 per cent, but this subject's scores were also the most reliable, for all this subject's trials were performed cooperatively, calmly, and methodically. This average percentage of difference was employed in computing the underhanded scores for the first and third subjects and for the left arm of the fourth; from the derived estimates, the adjusted average maximum values were calculated.

9. CONCLUSIONS

Bracing to avoid movement of the shoulder and observations of the changing angle of flexion at the elbow were employed in an attempt to minimize the chance of a decisive contribution from the scapular muscles, the significance of which is indicated by the 17.9 per cent increase (140 versus 165) in the maximum lift of the third subject in series D2 when the upper limb was fully extended. But the greater difficulty of making other requisite observations during the testing of the chimpanzee subjects and the greater difficulty of bracing the shoulders of the apes cause the writers to lack complete confidence that many of the chimpanzee scores are not somewhat higher due to the action of muscles of scapular origin; for example, among these muscles, the long head of triceps is relatively much heavier and has a longer scapular attachment than that of man (Edwards, 1965c). Also, the postulated frequent reliance upon elevation and retraction of the shoulder-joint (Hollinshead, 1951, pp. 91-97) would in part explain the difficulty experienced in attempting to teach the third and fifth subjects to pull at varied angles. On the other hand, flexion at the elbow was observed

Table 4

Weightlifting Performances of Chimpanzee Subjects

Subject	Maximum Lift (in pounds)	Adjusted Average Maximum Lift (in pounds)	Body-Weight (in pounds)	Average Maximum Lift in Pounds per Pound of Body-Weight	Average Brachial Cross-Sectional Area	Average Maximum Lift in Pounds per Square Centimeter
1	A 117	112.4	43.0	2.614	37.99	2.99
	B					
	C (140)					
	D					
2	A 110	105	44.5	2.360	38.51	2.73
	B 100					
	C					
	D					
3	A (110)	124.9	54.25	2.302	35.27	3.54
	B					
	C 130					
	D					
4	A 135	127.4	60.0	2.123	38.51	3.31
	B 115					
	C 135					
	D (100)					
5	A 270	260.0	98.0	2.653	73.06	3.56
	B 260					
	C 260					
	D 250					

A = Right, overhanded
 B = Right, underhanded
 C = Left, overhanded
 D = Left, underhanded

on many lifts and was in some cases described in the writer's notes immediately upon the tests' completion. Thus it is recorded that on 24 August the second subject "flexed arm but shoulder immobile" as for 3 seconds a 100-pound weight was lifted overhanded the entire range of handle-shaft motion; this lift was only 9.1 per cent less than the maximum. Likewise, all of the 1 September testing of the fifth subject was performed with meticulous bracing of the right shoulder with wooden blocks and with the left arm strapped to the left wall of the apparatus; yet overhanded and underhanded lifts were 265 and 250 pounds, only 1.9 and 3.8 per cent, respectively, less than previous maxima of 270 and 260 pounds. Furthermore, unlike the third human subject during series D2, the upper limbs of all the simian subjects were slightly flexed, and the muscular force required to maintain the slight flexion at the elbow during the pull is almost as demanding on the flexors of the elbow as their further contraction to produce additional flexion would be (Martin, 1921). Although any comparable testing program should seek to nullify shoulder movement more completely, perhaps by contoured braces of the shoulder and by training the chimpanzee subjects to pull at higher (though less optimal) angles of elbow flexion, the foregoing considerations indicate that any increase in these scores due to retraction at the shoulder is probably very moderate. It may therefore be concluded that with the satisfying of the recently analyzed thirty-seven criteria (incompletely satisfied in a few cases because of the limitations of available time) for adequate strength testing (Edwards, 1963a), both simian and human scores are quite accurately representative of the strength of the subjects tested, and interspecific comparisons are thus also valid.

The remarkable reliability of the chimpanzee scores accords with their validity. As indicated earlier in the paper, equivalent motivation was achieved through a widely varying assortment of positive and negative incentives, adjusted to the individual simian subject. It is of some interest to observe that only for the adult male was the primary motivation that of pleasing his human friends.

Intraspecific comparisons reveal the anticipated tendency for larger humans and chimpanzees to be stronger than smaller humans and chimpanzees (Edwards, 1965b). As would have been anticipated from studies of humans and from theoretical considerations, the maximal scores of chimpanzee females (second and third subjects) averaged less than those of males both in lift per unit of body-weight and in lift per unit of brachial cross-sectional area; but the number of subjects was too small and these ratios -- especially that of the force per unit of area, for which the third subject almost tied for highest score -- too inconsistent to be considered more than suggestive. The extent to which the scores of these second and third subjects averaged lower than those of the others might also be ascribed to the fact that they were the two youngest subjects. In any event, the fact that the highest scores both relative to brachial cross-sectional area and, despite the operation of geometrical similitude, relative to body-weight were those of the adult male chimpanzee may tend to some degree to confirm an anticipated correlation between strength and

both masculinity and maturity for these apes, as for humans; but firm conclusions are again precluded by the limited sample size.

Inspection of the data presented evinces the finding that, despite the fact that all of the human subjects were adult males but 40 per cent of the chimpanzees were females and 80 per cent were pre-adolescent or earliest-adolescent, chimpanzees are much stronger than humans per unit of body-weight. The outpulling by the largest chimpanzee of a human weight-lifter fully $2\frac{1}{2}$ times as large in body-weight seems especially noteworthy. The striking average ratio of superiority of chimpanzees to humans is 2.681:1 (2.410 to 0.899), approximating the ratio reported by Bauman (1923 and 1926). Much of this superiority is ascribable to geometrical similitude operating on the individuals of different body-size and to the proportionately heavier arms of the apes. For example, if a 64-pound chimpanzee, identical physiologically and in proportions to a 216-pound human, lifted 64 pounds, the human would lift 144 pounds, 33.3 per cent less than the 216 pounds which would have been lifted if strength were proportionate to the cube of a given dimension (or body-weight) instead of to the square of a given dimension (Edwards, 1963d). But general geometrical similitude provides only a very small factor in accounting for the 2.951:1 superiority of the adult male fifth subject.

As noted in a companion paper (Edwards, 1965b), unfortunately completed before the data here presented had been quantitatively analyzed, it is concluded that four to six other factors contribute significantly to the chimpanzee's marked superiority in strength. The most important of these other factors is the proportionately much greater massiveness of the chimpanzee upper limbs. Thus, per unit of brachial cross-sectional area, the chimpanzees manifest only a 41.1 percentage of average superiority over the humans (3.226 to 2.286). In fact, the ranges actually meet, but the immaturity and sexual heterogeneity of the chimpanzee subjects should again be emphasized, and it should be observed that only the proportionately weakest of the apes, a $4\frac{1}{2}$ -year-old female, demonstrated a ratio as low as the highest among the humans here tested.

Further comparisons between the two species reveal marked contrasts in the degree of overhanded superiority. The average amount by which the fatigue-adjusted (by 15 pounds for the second series maximum, as analyzed presently) underhanded scores were less than the overhanded scores for humans was 51.8 pounds (35, 62.5, 22.5, 35, 42.5, 75, 90 for the seven subjects, respectively). This average difference is equivalent to 27.7 per cent of the overhanded lifts, averaging 187.1 pounds, as compared with only some 7.85 per cent difference for chimpanzees, as discussed above. The writer has not yet had time to attempt a complete analysis of the presumed anatomical factors determining most of this contrast; but at least a small part of the contrast is surely due to the equivalent effort on all maximum lifts by the chimpanzees, while the human subjects were, in the opinion of the writer, more concerned about the maximum score for each arm than the overhanded-plus-underhanded average maximum.

The human maximum strength scores per square centimeter of brachial cross-sectional area manifest decidedly more interindividual variability than the chimpanzee scores, as shown by the contrasting ranges -- 1.70-2.77 versus 2.73-3.56 -- and by the contrasting average percentages by which the individual values differ from the mean values (2.286 and 3.226) in each group -- 13.9 versus 9.08. Thus, by the index here employed, human variability is some 53 per cent larger. Much of the lesser variability of the apes reflects more uniformity of exercise, but much of it is probably due to less genetic variability, as theorized in a previous paper (Edwards, 1965b) should be the case. Furthermore, chimpanzee intraindividual variability for maximum scores on different days also seems to be less than that indicated by the typical retest coefficient of correlation of .90 to .92 reported for humans (Hunsicker, 1955; Clarke, 1960), as was also theorized earlier on the basis of the chimpanzee's apparently greater ease of approaching strength maxima, presumably asymptotically (Edwards, 1963a, p. 14).

There is moderate indication of handedness in the humans tested, with higher scores for the preferred limb in all underhanded-plus-overhanded maximum scores except for the identical totals of the fourth subject, and with a 10.4-pound average superiority of the preferred arm in scores fatigue-adjusted for the first five subjects, as calculated below. Also, all seven humans exhibit slightly to moderately larger brachial girths for the preferred extremity. Insufficient time was available in the present study for observations on chimpanzee handedness per se. Since these apes are adapted to arboreal locomotion by varied climbing and brachiation, less handedness than in man would be anticipated theoretically (Edwards, 1963a and 1965b), and observational data procured by others confirm this expectation (Yerkes and Yerkes, 1929, p. 217). Some further confirmation of handedness to a lesser degree than in man is provided by the present study, as indicated by the generally greater similarity in chimpanzees of both contralateral strength scores and girths of the upper extremities. In the simian subjects -- two of whom manifested slightly larger right arm girths, two larger left arm girths, and one almost identical girths -- no clear correlation between the few contralateral strength scores and brachial girths is evident, but this is at least consistent with a more limited degree of handedness, although likely primarily reflecting the inadequate sample size. It may be somewhat significant to observe that in the subject most adequately tested, the largest, the right brachial cross-sectional area, roughly (as accurately as could be measured) 2.7 per cent larger, is associated with an average 3.9 per cent greater lift. In neither species does greater size and strength of one limb show direct genetic origin, however, since greater use of one arm results in its relative hypertrophy.

The two largest human subjects did not manifest any clear evidence of fatigue in the alteration of overhanded to underhanded sequences, for the overhanded superiority increased by 10 and by 20 pounds when preceded by the underhanded test. The lack of clearly manifested fatigue in these two humans may reflect better muscular endurance than that of the others, for one had for a year practiced weightlifting, while the other had during

the previous two months exercised his arms³ to an unusual extent; these two subjects were also likely the best motivated among the seven.

The five remaining humans apparently exhibited rather marked effects of fatigue by the lessening of the anticipated maximum score for the latter of the two series (overhanded and underhanded) for a given limb. The seven pairs with overhanded priority averaged a 57.1-pound overhanded superiority, but three pairs with underhanded priority averaged only an 18.3-pound overhanded superiority. It is estimated by approximate consensus of available data -- including some consideration of the inconsistent scores for the two largest subjects -- that the fatigue resulting from the first series for a given limb reduces the second series for that limb roughly 15 pounds.

In comparing the chimpanzee with the human performances, it seems evident that the proportionate superiority of chimpanzee endurance is much greater than that of strength, for many of the near-maximal pulls of the chimpanzees were made in much more rapid succession than those of the humans without apparent reductions in the scores achieved. Nevertheless, the fatigue factor was perhaps of slightly greater consequence in proportionate reductions of maximum scores achieved by the chimpanzees, for in addition to the briefer rest-intervals permitted between near-maximal pulls (in most cases a maximum of one minute for the apes compared to an average of more than two for the humans), much longer series of preliminary trials, primarily composed of lifts of less than 80 pounds not shown in the data previously presented, were mandatory for all but the largest chimpanzee subject. Furthermore, many successive series were generally employed for the apes, in many cases during four to six hours without a single very lengthy rest-interval.

There might also have been anticipated some evidence of a general fatigue factor in the human tests. Comparison of intralimb fatigue-adjusted (15 pounds for the first five men) and interlimb handedness-adjusted (10 pounds for all humans) scores reveals that the first limb to be tested scored higher than was anticipated by an average superiority of only 1.79 pounds over the contralateral limb for the fourteen pairs compared. It should be considered, however, that even should a larger series also fail to yield statistically significant results, a general fatigue factor would not necessarily be invalidated, for there may also be psychological, epinephrine-associated, and/or other compensating factors. The chimpanzee test results failed to manifest any such correlation, as would have been anticipated in view of the smaller sample of comparative scores and the better endurance of the apes.

Through the study here presented, it is now possible to state the approximate strength of chimpanzees relative to humans. Chimpanzees are two to three times as strong as humans per unit of body-weight and, at least for the brachial flexors, roughly 50 per cent stronger than humans per unit of cross-sectional muscle area, sex and age being equivalent.

REFERENCES

- Bauman, J. E. 1923. "The Strength of the Chimpanzee and the Orang." Scientific Monthly, Vol. 16, pp. 432-439.
- Bauman, J. E. 1926. "Observations on the Strength of the Chimpanzee and its Implications." Journal of Mammalogy, Vol. 7, pp. 1-9.
- Clarke, D. H. 1960. "Correlation between the Strength/Mass Ratio and the Speed of an Arm Movement." Research Quarterly, Vol. 31, pp. 570-574.
- Cureton, T. K. 1947. Physical Fitness Appraisal and Guidance. C. V. Mosby Co., St. Louis.
- Edwards, W. E. 1963a. "The Testing of Chimpanzee Strength Prior to 1961: Methods and Results." Technical Documentary Reports, No. ARL-TDR-63-22, 6571st Aeromedical Research Laboratory, Holloman Air Force Base, New Mexico.
- Edwards, W. E. 1965b. "Factors in the Superiority of Chimpanzee over Human Strength." Technical Documentary Reports, 6571st Aeromedical Research Laboratory, Holloman Air Force Base, New Mexico.
- Edwards, W. E. 1965c. "The Musculoskeletal Anatomy of the Thorax and Brachium of an Adult Female Chimpanzee." Technical Documentary Reports, 6571st Aeromedical Research Laboratory, Holloman Air Force Base, New Mexico.
- Edwards, W. E. 1963d. "The Relationships of Human Size to Strength." Technical Documentary Reports, No. ARL-TDR-63-19, 6571st Aeromedical Research Laboratory, Holloman Air Force Base, New Mexico.
- Finch, G. 1943. "The Bodily Strength of Chimpanzees." Journal of Mammalogy, Vol. 24, pp. 224-228.
- Hollinshead, W. H. 1951. Functional Anatomy of the Limbs and Back. W. B. Saunders Company, Philadelphia.
- Hunsicker, P. A. 1955. "Arm Strength at Selected Degrees of Elbow Flexion." Technical Report 54-548, Wright Air Development Center, Wright-Patterson Air Force Base, Ohio.
- Martin, E. G. 1921. "Tests of Muscular Efficiency." Physiological Reviews, Vol. 1, pp. 454-475.
- Yerkes, R. M. and Yerkes, A. W. 1929. The Great Apes: A Study of Anthro-poid Life. Yale University Press, New Haven.
- Wilkie, D. R. 1950. "The Relationship between Force and Velocity in Human Muscle." Journal of Physiology, Vol. 110, pp. 249-280.

DISTRIBUTION

AFSC (SCTB) Andrews AFB Wash, DC 20331	6	Academy Library-DFS LB United States Air Force Academy Colorado 80840	2
Hq, USAF, Science Division (AFRSTA) Dir/Science and Technology, DCS/R&D Wash, DC 20330	1	American Institute of Aero- nautics and Astronautics 750 Third Avenue New York, NY 10017	1
AMD ATTN: Chief Scientist Brooks AFB, Texas 78236	1	Boeing Airplane Company Aero-Space Division ATTN: Ruth E. Peerenboon, Process Supervisor P. O. Box 3707 Seattle, Wash 98124	1
AMD (AMAP) Brooks AFB, Texas 78236	10	Redstone Scientific Information Center ATTN: Chief, Document Section U.S. Army Missile Command Redstone Arsenal, Ala 35809	5
DDC (TIAAS) Cameron Station Alexandria, Va 22314	20	British Liaison Office Army Missile Test and Evalu- ation Directorate White Sands Missile Range New Mexico 88002	1
AFETR Tech Library (MU-135) Patrick AFB, Fla 32922	1	Defence Research Member Canadian Joint Staff Director of Biosciences Research 2450 Massachusetts Ave., N.W. Wash, DC 20008	1
APGC (PGBAP-1) Eglin AFB, Fla 32542	1	USAF School of Aerospace Medicine ATTN: Aeromedical Library Brooks AFB, Texas 78236	1
ESD (ESTI) L.G. Hanscom Field Bedford, Mass 01730	2	Life Sciences Dept., Code N300 U.S. Naval Missile Center Point Mugu, Calif 93041	1
Air University Library Maxwell AFB, Ala. 36100	1		
Central Intelligence Agency ATTN: OCR/DD-Standard Distribution Wash, DC 20505	2		

Aviation Medical Acceleration Laboratory ATTN: AMAL Library Johnsville, Pa 18974	2	Medical Records Section Room 325 Division of Medical Sciences National Academy of Sciences National Research Council 2101 Constitution Avenue NW Wash, DC 20418	1
Librarian C.A.R.I. F.A.A. P.O. Box 1082 Oklahoma City, Okla 73101	1	Aviation Safety Engineering Research A Division of Flight Safety Foundation 2871 Sky Harbor Blvd Sky Harbor Airport Phoenix, Arizona 85034	1
Director U.S. Naval Research Laboratory (Code 5360) Wash, DC 20390	2	NASA Langley Research Center Langley Station ATTN: Librarian MS185 Hampton, Va 23365	3
Biomedical Library Center for Health Sciences University of California Los Angeles, Calif 90024	1	Archive-Library Dept American Medical Assoc 535 N. Dearborn St., Chicago, Ill 60610	1
Technical Reference Library U.S. Naval Medical Research Institute Bethesda, Md 20014	1	Information Officer USAFE French Liaison Office Det 0100, 7260th Support Group APO New York, NY 09686	1
Director Walter Reed Army Institute of Research ATTN: Neuropsychiatry Division Wash, DC 20012	1	The Decker Corp 45 Monument Road Bala-Cynwyd, Pa 19004	1
Commanding Officer ATTN: Code 373 U.S. Naval School of Aviation Medicine U.S. Naval Aviation Medical Center Pensacola, Fla 32512	2	Douglas Aircraft Co Advance Biotechnology Dept Missile and Space Systems Div Santa Monica, Calif 90406	1
Dr. W.M. Helvey 0/55-60 B/151 Lockheed Missiles and Space Co P.C. Box 504 Sunnyvale, Calif 94088	1	Literature Acquisition Dept Biological Abstracts 3815 Walnut St Philadelphia, Pa 19104	1

The Lovelace Foundation Dept of Aerospace Medicine and Bioastronautics 5200 Gibson Blvd., S.E. Albuquerque, NMex., 87108	1	Mr. Elliott H. Morse Librarian Library of the College of Physicians of Philadelphia 19 South 22nd Street Philadelphia, Pa 19103	1
Dr. William D. Thompson Department of Psychology Baylor University Waco, Texas	1	Dr. Leon H. Schmidt, Director National Center for Primate Biology University of California Davis, California 95616	1
Dr. Deets Pickett 8505 Lee Blvd Leawood, Kansas 66206	1	Del Electronics Corp Research & Development Dept ATTN: Leonard R. Solon, Ph.D. 250 E. Sanford Blvd., Mount Vernon, New York	1
Dr. Walter J. Frajola M-351 Starling-Loving Hall Ohio State University Columbus 10, Ohio 43210	1	Civil Aeromedical Research Institute Federal Aviation Agency Aeronautical Center ATTN: John N. Ice, AM-119 Editor, Acceleration Project P.O. Box 1082 Oklahoma City, Oklahoma 73101	1
Dr. R.D. Gafford Project Manager, Space Engineering Beckman Instruments, Inc 2500 Harbor Blvd Fullerton, Calif 92632	1	Director Flyvemedicinsk Institute Copenhagen, Denmark	1
Commanding Officer U.S. Naval Medical Field Research Laboratory Camp Lejeune, NC ATTN: Library	1	National Aeronautics and Space Administration Scientific and Technical Infor- mation Facility ATTN: NASA Representative Bethesda, Maryland 20014	2
Aerospace Medicine Bibliography Project Science and Technology Division Library of Congress Wash, DC 20540	1	Human Research Branch (Code RBH) OART, NASA Headquarters Wash, DC 20546	2

Hq, U. S. Army Medical R&D 1
Command
ATTN: Chief, Behavioral Sciences
Research Branch
Room 2608, Main Navy Bldg.,
Wash, D.C. 20315

D. L. Patterson 1
School of Veterinary Medicine
University of Pennsylvania
Philadelphia, Pa

Dr. Clement A. Stone, Director 1
Merck Institute of Therapeutic
Research
West Point, Pa 19486

Dean, College of Veterinary 2
Medicine
Iowa State University
Ames, Iowa 50010

Dean 2
School of Veterinary Medicine
Ohio State University
1900 Coffey Road
Columbus 8, Ohio 43210

C. J. Marshall Memorial Library 2
School of Veterinary Medicine
University of Pennsylvania
3800 Spruce Street
Philadelphia, Pa 19104

Dr. W. R. Pritchard 2
Dean, School of Veterinary
Medicine
University of California
Davis, Calif 95616

Dean C. A. Brandly 2
College of Veterinary Medicine
University of Illinois
Urbana, Ill 61803

Dean 2
School of Veterinary Medicine
Oklahoma State University
Stillwater, Okla

Biosciences Division 1
Eoar Shell Bldg
47 Cantersteen
Brussels, Belgium

Dr. J. E. Green, Dean 2
School of Veterinary Medicine
Auburn University
Auburn, Ala 36830

Library, New York State 2
Veterinary College at
Cornell University
Ithaca, NY 14850

Dean, Ralph L. Ketchell 2
College of Veterinary Medicine
Kansas State University
Manhattan, Kansas 66504

Dean 2
College of Veterinary Medicine
Michigan State University
East Lansing, Mich

Dean 2
School of Veterinary Medicine
University of Georgia
Athens, Ga

Dr. Robert L. Hamlin 1
1900 Coffey Road
Columbus 10, Ohio 43210

Librarian 1
Department of Health, Education
and Welfare
Public Health Service
Occupational Health Research and
Training Facility
Bureau of State Services
1014 Broadway
Cincinnati, Ohio 45202

Library 1
National Institutes of Health
Building 10, Room 5N115
ATTN: Acquisitions Unit
Bethesda, Maryland 20014

Major Joung Kook Park 1
ROKAF (MC)
Office of Surgeon General
HQ ROKAF, Seoul, Korea

Library 1
Oregon Regional Primate
Research Center
505 N.W. 185th Avenue
Beaverton, Oregon 97005

Dean, School of Veterinary Medicine
Washington State University 1
Pullman, Washington

Local

MDNH	1
NLO	1
ARGS	80
RRRT	3
RRRS	1

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1 ORIGINATING ACTIVITY (Corporate author) William E. Edwards		2a REPORT SECURITY CLASSIFICATION Unclassified	
		2b GROUP	
3 REPORT TITLE STUDY OF MONKEY, APE, AND HUMAN MORPHOLOGY AND PHYSIOLOGY RELATING TO STRENGTH AND ENDURANCE - PHASE IX: THE THE STRENGTH TESTING OF FIVE CHIMPANZEE AND SEVEN HUMAN SUB- JECTS			
4 DESCRIPTIVE NOTES (Type of report and inclusive dates) Final report			
5 AUTHOR(S) (Last name, first name, initial) Edwards, William E. and Clarke, Thomas Erskine			
6 REPORT DATE August 1965		7a TOTAL NO OF PAGES 21	7b NO OF REFS 14
8a CONTRACT OR GRANT NO. AF 29(600)-3466 b. PROJECT NO. 6892 c Task No. 689201 d		9a ORIGINATOR'S REPORT NUMBER(S) ARL-TR-65-15 9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10 AVAILABILITY/LIMITATION NOTICES			
11 SUPPLEMENTARY NOTES		12 SPONSORING MILITARY ACTIVITY	
13 ABSTRACT Five chimpanzees (two immature of each sex and one adult male) were trained for testing of elbow flexion strength. Resulting scores were compared with those of seven young (20 to 37 years) adult human males. The apes manifested a 2-1/2 fold superiority by body-weight and, sex and age equivalent, an appreciable superiority by brachial cross-sectional area.			

14

KEY WORDS

Chimpanzees
Humans
Strength
Endurance
Elbow flexion

LINK A

ROLE

WT

LINK B

ROLE

WT

LINK C

ROLE

WT

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.